

OROLONE PASTE®

Triamcinolone Acetonide 0.1%

TARGET THE PROBLEM:

Provide the solution

PRESENTATION:

Adhesive ointment; 0.1% triamcinolone acetonide, dental paste with additional flavour.

PRESENTATION:

Adhesive ointment; 0.1% triamcinolone acetonide, dental paste with flavour.

PHARMACOLOGICAL CLASS:

Corticosteroid.

Actions:

Triamcinolone acetonide is a synthetic corticosteroid which possesses anti-inflammatory anti-pruritic, and anti-allergic action which provide prompt relief of oral tenderness, pain, inflammation. Triamcinolone is used to reduce swelling and ulcers in the mouth. The OROLONE paste is medication for applying to oral surfaces. This adhesive maintains the medication in close contact with the lesion.

INDICATIONS:

OROLONE paste is indicated for adjunctive treatment and for the temporary relief of symptoms associated with oral inflammatory lesions.

OROLONE paste is used to treat inflamed mouth sores and mouth sores caused by injury. This medication is used for the temporary relief of symptoms from mouth sores due to injury. It is applied directly to the affected areas of the mouth. It acts by decreasing inflammation and protecting the damaged area from further injury.

This medication is used for the temporary relief of symptoms from mouth sores due to injury. This drug is in the form of dental paste that allows it to stick to the inside of the mouth/cheeks/gums. It works by reducing the swelling, itching and pain that can occur with mouth sores.

DOSAGE AND ADMINISTRATION:

Apply a small amount as per the size of lesion to the affected area. A larger quantity may be required for coverage of some lesions. The preparation should be applied at bedtime to permit steroid contact with the lesion throughout the night. Depending on the severity of symptoms, it may be necessary to apply the preparation two to three times a day, preferably after meals. If significant improvement has not occurred in seven days, further investigation is advisable.

ADULT:

Press (do not rub) thin film onto lesion at bedtime and, if needed, 2-3 times daily after meals. Reevaluate if no improvement in 7 days.

C Apply this medication 2 or 3 times daily, preferably after meals. At least one application should be at bedtime so that the medication can be in contact with the lesion overnight.

CHILDREN:

Not recommended.

CONTRAINDICATIONS:

This preparation is contra-indicated in patients with a history of hypersensitivity to Triamcinolone acetonide. Because it contains a corticosteroid, the preparation is contraindicated in the presence of fungal, viral, or bacterial infections of the mouth or throat.

PRECAUTIONS:

Use during Pregnancy

Safe use of this product during pregnancy has not been; therefore it should not be used in women of child bearing potential and particularly during early pregnancy unless in the judgement of the physician or dentist the potential benefits outweigh the possible hazards.

SIDE EFFECTS

Local Minor Irritation and burning

Prolonged administration may elicit the adverse reactions known

ओरोलोन

to occur with systemic steroid preparations; for example adrenal suppression, alteration of glucose metabolism, protein catabolism, peptic ulcer activations, and others. These are usually reversible and disappear when the hormone is discontinued.

Manufactured by:

Deurali-Janta Pharmaceuticals Pvt. Ltd.

679, Budhanilkantha Sadak, Sansban - 03 | G.P.O. Box: 4239, Kathmandu, Nepal
Tel: +977-01-4018777 | E-mail: mplaning@deuralijanta.com

Website: www.deuralijanta.com

© Registered Trade Mark

01 pi OPD 20

