

DIGIROLFIN 5% डिजिरोलफिन ५%

AMOROLFINE 5% W/V MEDICATED NAIL LACQUER

INFORMATION FOR THE USER

DIGIROLFIN 5% w/v Medicated Nail Lacquer

Amorolfine

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

1. Contents of the pack and other information

What DIGIROLFIN 5% contains

The active substance is amorolfine. DIGIROLFIN 5% contains 50 mg/ml (5%) of amorolfine (equivalent to 6.4% amorolfine hydrochloride).

2. What DIGIROLFIN 5% is and what it is used for ?

DIGIROLFIN 5% is used to treat fungal infections of the nails.

DIGIROLFIN 5% contains the active ingredient amorolfine (as the hydrochloride), which belongs to a group of medicines known as antifungals. It kills a wide variety of fungi that can cause nail infections.

3. What you need to know before you use DIGIROLFIN 5% ?

Do not use DIGIROLFIN 5% if you are:

- Allergic (hypersensitive) to amorolfine or any of the other ingredients of DIGIROLFIN 5%.

Warnings and precautions

This medicine can cause allergic reactions, some can be serious. If this happens, stop applying the product, immediately remove the product with a nail varnish remover or the cleaning swabs and seek medical advice. The product must not be reapplied. You must get urgent medical help if you have any of the following symptoms:

- You have difficulty breathing
- Your face, lips, tongue or throat swell
- Your skin develops a severe rash

Talk to your doctor or pharmacist before using DIGIROLFIN 5%.

The lacquer should not be applied on the skin around the nail.

- Avoid the lacquer coming into contact with the eyes, ears or mucous membranes (e.g. mouth and nostrils).
- Do not inhale.
- Wear impermeable gloves when using organic solvents to stop the nail lacquer from being removed.

Using other medicines

There are no known interactions with other medicines.

Using other nail products

Artificial nails should not be used while using DIGIROLFIN 5%.

Cosmetic nail varnish can be used, but you should wait at least 10 minutes after applying DIGIROLFIN 5% before painting your nails.

The nail varnish should be carefully removed before repeat application of DIGIROLFIN 5%.

Pregnancy and breast-feeding

You should not use DIGIROLFIN 5% and tell your doctor if you are pregnant, planning to become pregnant or are breast-feeding. Your doctor will then decide whether you should use DIGIROLFIN 5%.

Ask your doctor or pharmacist for advice before taking any medicine.

4. How to use DIGIROLFIN 5%

Always use this medicine exactly as your doctor had told you. Check with your doctor or pharmacist if you are not sure. DIGIROLFIN 5% should be applied to the affected finger or toe nails once or twice a week exactly as directed by your doctor.

Instructions for use

Step 1. File the nail

Before the first application, file down the infected areas of nail, including the nail surface, as much as possible.

CAUTION: Do not use nail files used for infected nails on healthy nails, otherwise you may spread the infection.



Step 2. Clean the nail

Use swab to clean the nail surface.

Repeat steps 1 and 2 for each affected nail.



Step 3. Take some lacquer from the bottle

Dip the applicator into the bottle of nail lacquer. The lacquer must not be wiped off on the edge of the bottle before it is applied.



Step 4. Apply the lacquer

Apply the nail lacquer evenly over the entire surface of the nail.

Repeat steps 3 and 4 for each affected nail.



Step 5. Allow to dry

Let the treated nail(s) dry for approximately 3 minutes. Wait at least 10 minutes before applying cosmetic nail varnish.



DIGIROLFIN 5% डिजिरोलफिन ५%

Step 6. Clean the applicator

The applicator provided is re-usable. However, it is important to clean it thoroughly after completing each treatment procedure, using the swab. Avoid touching newly treated nails with the swab. Close the nail lacquer bottle tightly. Dispose of the swab carefully as it is inflammable.



- Before using DIGIROLFIN 5% again, first remove the old lacquer and any nail varnish from your nails using a swab, then file down the nails again if necessary.
- Re-apply the lacquer as described above.
- When dry the nail lacquer is unaffected by soap and water, so you may wash your hands and feet as normal. If you need to use chemicals such as paint thinners or white spirit, rubber or other impermeable (water-proof) gloves should be worn to protect the lacquer on your fingernails.
- It is important to carry on using DIGIROLFIN 5% until the infection has cleared and healthy nails have grown back. This usually takes 6 months for finger nails and 9 to 12 months for toe nails. Your doctor will probably check how your treatment is progressing every 3 months or so.

If you get DIGIROLFIN 5% in your eyes or ears

If you get DIGIROLFIN 5% in your eyes or ears wash out with water immediately and contact your doctor, pharmacist or nearest hospital straight away.

If you accidentally swallow DIGIROLFIN 5%

If you, or anyone else, accidentally swallows the lacquer contact your doctor, pharmacist or nearest hospital straight away.

If you forget to use DIGIROLFIN 5%

Don't worry if you forget to use DIGIROLFIN 5% at the right time. When you remember, start using the product again, in the same way as before.

If you stop using DIGIROLFIN 5%

Do not stop using DIGIROLFIN 5% before your doctor tells you to or your infection could come back.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

5. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. In addition to the beneficial effects of DIGIROLFIN 5%, it is possible that unwanted effects will occur during treatment, even when it is used as directed.

Rare: may affect up to 1 in 1,000 people

- Nail damages, nail discoloration, fragile or brittle nails.

Very rare: may affect up to 1 in 10,000 people

- Skin burning sensation.

Not known: frequency cannot be estimated from the available data

- Systemic allergic reaction (a serious allergic reaction that can be associated with swelling of the face, lips, tongue or throat, difficulty breathing and/or a severe skin rash)
- Redness, itching, hives, blister, allergic skin reaction

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via www.deuralijata.com. By reporting side effects you can help provide more information on the safety of this medicine.

6. How to store DIGIROLFIN 5% ?

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the pack. The expiry date refers to the last day of that month.
- Keep the pack away from heat (store below 25°C) and keep the bottle tightly closed after use. Do not throw away any medicines via waste-water or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

7. What DIGIROLFIN 5% looks like and contents of the pack ?

The pack is available in size of 2.5 ml bottle.

Each 2.5 ml pack consists of 1 bottle filled with DIGIROLFIN 5% nail lacquer and spatula integrated in the cap.



Manufactured by:

Deurali-Janta Pharmaceuticals Pvt. Ltd.

679, Budhanilkantha Sadak, Bansbari - 03 | G.P.O. Box: 4239,

Kathmandu, Nepal Tel: +977-01- 4018777

E-mail: mplanning@deuralijanta.com, Website: www.deuralijanta.com

02 pi DRS 24

